

## **Impact of Self-Help Psychoeducational Intervention on Management of Emotional Eating among Obese Individuals: A Randomized Controlled Trial**

**Nimra Anjum, Sajida Naz, Sadia Aleem, and Sana Kiaani**

Fatima Jinnah Women University

**. Ghulam Saqulain**

Capital Hospital, Islamabad

With a high prevalence of emotional eating affecting young adults in Pakistan, leading to obesity and other health problems there is urgent need of addressing this issue to prevent severe mental, physical, and social health impairments; and potential benefit of psycho-educational intervention, the current research was conducted to evaluate the effectiveness of a psycho educational program on overweight/ obese young adults by targeting emotional eating, psychological distress, social support and quality of life, using a two-arm, single-blinded, randomized control trial at Fatima Jinnah Women University, Rawalpindi from Nov 2023 to Jan 2024. Psycho-educational Weight Reduction Program can widely be used as a method to improve functioning in young adults having extra build weight and also reduce emotional eating. It can be presumed that changes of eating behaviors, emotional functioning and body image may be imperative in weight management program Study included two phases with a sample of 550 students recruited using convenience sampling with cross-section study design in Phase-I and sample screened for high emotional eating (score >75) and Body Mass Index (25-30 Kg/m). Fourteen eligible students entered Phase II (Intervention) and randomly assigned into the Experimental (self-help psycho-educational weight loss intervention for 12 weeks (2-sessions/ week) and Control groups. Pre and Post-intervention results revealed that the experimental group had significantly lower scores ( $P < 0.001$ ) for Generalized Anxiety Disorder, Emotional Eating Scale, Patient Health Questionnaire and Discouragement

---

Nimra Anjum, Sajida Naz, Sadia Aleem, and Sana Kiaani, Department of Behavioral Sciences, Fatima Jinnah Women University, Pakistan.

Ghulam Saqulain, Head of Department & Professor, Department of Otorhinolaryngology & Head and Neck Surgery; Capital Hospital, Islamabad, Pakistan.

Correspondence concerning this article should be addressed to Ghulam Saqulain, Head of Department & Professor, Department of Otorhinolaryngology & Head and Neck Surgery; Capital Hospital, Islamabad, Pakistan. Email: [ghulam\\_saqulain@yahoo.com](mailto:ghulam_saqulain@yahoo.com)

subscale, while significantly higher scores were noted for Physical Component Summary(PCS), Mental Component Summary(MCS) and Encouragement subscale, compared to Control Group. Hence, Self-help psycho educational program for management of emotional eating has significant effects on young adults having obesity in terms of reducing emotional eating, psychological distress and improving social support and life quality.

*Keywords.* Emotional eating, Body Mass Index, Psychological, Quality of life, and self-help psycho-educational programs

Spread of COVID-19 starting from 2020 emerged as a significant public health concern, profoundly disturbing daily routines worldwide and exerting substantial influences on both physical and mental well-being (Holmes et al., 2020). This was followed by regional lockdowns, as a means to limit its transmission (Ghebreyesus, 2020). Social isolation and quarantine protocols adversely impacted on individuals' physical and mental well-being. Mental health repercussions of COVID-19 pandemic were experienced by the entire populace, with a more pronounced effect on females, young adults, healthcare professionals, and individuals with lower socioeconomic status. This is due to factors such as women's increased responsibilities in childcare and employment within sectors disproportionately affected by the pandemic (Niedziedz, 2021).

Emotional Eating is considered Obeseogenic with contribution in achieving an increase in. It commonly results from negative emotions (Kontinen, 2020). Psychosomatic theory explains the emotional eating concept. Individuals with emotional eating fail to see the difference between starvation and physiological condition related to negative emotion. Historically, emotional eating is related to tendency of increasing diet intake because of altered emotions which may be in form of individual's state of being angry, depressed, furious, nervous or lone (Güneşer & Him, 2021).

According to World Health Organization, the prevalence of this emotional state in the local context is considered 58.1 percent, however 43.9% suffer obesity (Coronavirus Disease 2019 (COVID-19): Situation Report - 64 (24 March 2020) - World, 2020). This indicates Pakistan has to cater to the factors which predict obesity, and essentially catering to the major factor ie., Emotional Eating. Pre-pandemic, 38% adults admitted to overeating while during this period it rose (47%) (Emotional Eating on the Rise During the Pandemic, 2020).

Etiology of Emotional Eating is not known. But, supposedly it can result from a combination of issues related to environment, psychology

and biology with their basis on a “bio-psycho-social” models. In other words it can be phrased as “Gun is loaded by the genetic makeup and its trigger pulled by the environment (Galmiche et al., 2019).

The potential benefits of using a psycho-educational weight reduction program is enhancement of functioning of youth with weight gain and increase emotional eating. Hence, it is suggested that variations in behavior of eating, emotions, self-image are crucial in programs which aim to manage weight (Lattimore, 2019). The self-help approach is particularly useful for individuals who cannot access obesity treatment centers or psychological assistance in their area (Levin et al., 2018). The WHO and the European Association for the study of overweightness recommended that finding modern methods to help young adults and develop novel interventions that should be efficient, operative and extra suitable to requirements (McGowan 2016; World Health Organization 2015). These commendations were based on the statistic that: (I) the spread of obesity was continuously snowballing and the price of treatment continuously on a rise, (II) Few studies have showed that psychological factors are intricate in the maintenance, growth and handling of obesity (McGowan, 2016; World Health Organization 2015). However, concluded from previous research studies indicate that traditional treatment for obesity should be supplemented by psycho-education based on web that would be more effective in treating obesity related issues such as Emotional Eating. Research is required to highlight connections between emotional eating and weight gain and psychological distress

With a high prevalence of emotional eating affecting young adults in Pakistan ( [Coronavirus Disease 2019 \(COVID-19\)- Situation Report - 64 \(24 March 2020\) - World, 2020](#)), leading to obesity and other health problems there is urgent need of addressing this issue to prevent severe mental, physical, and social health impairments; and potential benefit of psycho-educational intervention (Lattimore, 2019), the current research was conducted to look into the effectiveness of a psychoeducational program on overweight/ obese young adults by targeting emotional eating, social support, psychological distress and quality of life.

This study has significant since it might have implications for clinical management of overweight/ obese emotional eaters and for research.

### **Objective**

1. To assess the efficacy of psycho-education self-help behavioral intervention in reducing Emotional Eating among overweight/obese young adults

2. To examine changes in psychological distress levels following the intervention
3. To evaluate improvements in perceived social support after the intervention
4. To determine changes in quality of life among participants post-intervention

### **Hypothesis**

Bases on the objective to evaluate the effectiveness of the psychoeducational program, the following hypotheses were formulated:

1. The program will significantly reduce emotional eating among overweight/obese young adults
2. The program will significantly reduce psychological distress
3. The program will significantly reduce perceived social support
4. The program will significantly enhance overall quality of life

### **Method**

Current study utilized Randomized Control Trial (RCT) design at Fatima Jinnah Women University (FJWU) Rawalpindi, after ethical approval from FJWU Ethics Committee approval vide reference number FJWU/ EC/2023/59 dated September 13, 2023 and RCT registration number NCT06081023. Study was carried out over a period 3 months starting from November 1, 2023 to January 31, 2024.

#### ***Phases and Participants***

Although the study was conducted in two phases, the data were analyzed and presented collectively to provide a comprehensive, objective-based interpretation rather than a phase wise breakdown, in line with the study's cross-sectional design.

Phase I: A Cross-Sectional Survey design using convenience sampling recruited a sample of 550 students from Foundation University, Arid agricultural University, Riphah International University, Barani Institute and National University of Modern Languages Rawalpindi/ Islamabad. Sample of 558 was calculated utilizing Raosoft calculator using a confidence level = 95%, 5% margin of error with a population of 3500. However eight participants refused

consent and were excluded. The sample of  $N = 550$  was screened for high Emotional Eating Scores and Body Mass Index (BMI). Participants with scores exceeding 75 were categorized as 'Emotional Eaters', while those scoring below 75 were classified as 'non-emotional eaters.' Of the total of 550 participants, it was determined that 50 individuals (10%) self-reported high levels of emotional eating. Moreover, the examination of Body Mass Index (BMI) data revealed that approximately 88 participants (16%) were categorized as overweight, on the BMI calculator. Overall screening revealed  $N = 15$  (2.73%) students who had high Emotional Eating Scores and BMI 25-30 Kg/m (Table 1) which was our sample of study in Phase II.

**Table 1:** *Assessment of Emotional Eating and Body Mass Index (N = 550)*

Variable	Group	Emotional Eating		
		Low	High	Total
Body Mass Index	Normal (18.5-25 kg/m)	427 (77.64)	35 (6.36)	462 (84)
	Overweight (25-30 kg/m)	73 (13.27)	15 (2.73)	88 (16)
	Total	500 (90.91)	50 (9.09)	550

These screened individuals were then informed about the opportunity to voluntarily participate in the second phase of the study. Sample comprised participants with variation of age from 19-35 with a mean age of 22 years for males and 21 years for females. These included 261(23.5%) females and 289 (52.5%), males.

Phase II: Of the sample screened out for the phase II i.e.,  $N = 15$ , one participant refused consent to undergo Phase II, while,  $N = 14$  expressed their commitment to the study for a period of 3 months and were randomly assigned to either the control or experimental group, constituting Phase II of the study which comprised a psycho-educational weight loss intervention utilizing a Randomized Control Trial. Each participant was provided with a volunteer participation form, and their involvement in the interventional study was acknowledged, including their right to withdraw from the study if they wished to do so.

## **Measures**

Evaluation tools included Emotional Eating Scale (EES) (Tanofsky-Kraff et al., 2007) which is a 25 item reliable tool that measures emotional eating using a 5 point Likert scale; Generalized Anxiety Disorder (GAD) Scale (Spitzer et al., 2006) measuring Anxiety using 7 items that are measured using a 4 point likert scale; Patient Health Questionnaire (PHQ) (Kroenke et al., 2001) measuring depression using 9-items with 4 point likert scale; Social Support for Eating Habits (SSEH) (Sallis et al., 1987) measuring social support using 10 items with 5 point likert scale; and Short Form 36 (SF-36) (Ware & Sherbourne, 1992) measuring general health using 36 items.

## ***Intervention***

Following pre-intervention assessment, the intervention, known as Self-Help, was then implemented exclusively for the experimental group. In contrast, the control group participants were provided with an audio clip that guided them through a muscle relaxing technique to alleviate psychological distress, but no other behavioral techniques were applied. Intervention was instituted for a period of 12 weeks. After the intervention, assessment was carried out for determination of effectiveness of the Self-Help intervention.

Reduction of weight using psycho-educational strategy can be utilized to enhance functions among youth who are obese and to cut down overeating associated with psychological issues. It is assumed that alteration in eating habit, emotional state and self-image is essential for managing weight in weight reduction programs (Montesi et al., 2016; Teixeira et al., 2015). The fact behind might be that improvement in mental health could have an enhanced role, raising motivation and self-control (Teixeira et al., 2015). Self-help strategy could help individuals living far-away from obesity-handling-centers. Further, web-based management may be a reasonable option for such individuals living far-away from these centers or in conditions like Covid-19 pandemic.

## ***Data Collection Procedure***

Data was collected pre-intervention and post-intervention by the researcher using the measures described above.

### *Statistical Analysis*

Data analysis for this study was performed using SPSS version 26. An independent t-test analysis was performed to juxtapose participants from distinct groups, focusing on emotional eating, anxiety, depression, social support, and quality of life for comparative assessment. *t*-test will be used to see difference between groups.  $P < 0.05$  was taken as significant value.

## **Results**

### **Demographic characteristics**

A randomized control trial was conducted with a cohort 14 cases with 07 cases Group A (Experimental Group) and 07 in Group B (control group). Out of the 14 participants screened, 6(42.86) were males and 8(57.14) females. The age range of most participants was 21-25 years with most being overweight 12(85.71%). No significant difference was noted among experimental and control groups with  $P > 0.05$  as regards age, gender and education level (Table 2).

**Table 2:** *Demographic Variables Versus Frequency of Participants Among Study Groups in Phase II: Cross Tabulation (N = 14)*

Variables	Category	Group		Total	p-Value
		Experimental (n = 7)	Control (n = 7)		
Gender	Male	3	3	6	0.704
	Female	4	4	8	
Age	20	2	0	2	0.502
	21	1	3	4	
	22	1	1	2	
	23	1	1	2	
	24	1	2	3	
Body Mass Index	Over weight (25-29.9)	5	7	12	0.462
	Obese (above 30)	2	0	2	

*Note.* PHQ = Patient Health Questionnaire, PCS = Physical Component Summary, MCS = Mental Component Summar, GAD = Generalized Anxiety Disorder).

**Table 3:** Mean Differences in Predictor Variables Between Pre and Post-Intervention Group (N = 14)

Variables	Pre-Intervention	Post-Intervention
	Mean±SD	Mean±SD
Emotional Eating Scale	87.07±7.28	64.79±26.91
t,p-value	3.495, 0.004	
PCS	49.00±8.88	57.29±14.43
t,p-value	-2.312, 0.038	
MCS	41.00±5.99	48.71±12.00
t,p-value	-2.209, 0.046	
GAD	14.86±2.85	8.29±8.11
t,p-value	3.192, 0.007	
PHQ	18.64±3.15	9.86±9.83
t,p-value	3.272, 0.006	
Encouragement	13.79±3.51	16.14±4.26
t,p-value	-1.612, 0.131	
Discouragement	16.79±1.63	14.79±6.13
t,p-value	1.05, 0.313	

Note. PHQ = Patient Health Questionnaire, GAD = Generalized Anxiety Disorder, MCS = Mental Component Summary, PCS = Physical Component Summary.

Results of the current study (Table 3) for scores of Emotional Eating Scale (EES), Physical Component Subscale (PCS), Mental Component Scale (MCS), Generalized Anxiety Disorder (GAD), Physical Health Questionnaire (PHQ), Encouragement and Discouragement for pre-intervention versus post-intervention results revealed significantly improved results for EES, PCS, MCS, GAD, PHQ, encouragement with  $p < 0.05$ ; while no significant ( $p = 0.313$ ) difference was noted for discouragement.

**Table 4:** Mean Differences in Predictor Variables Between Control and Experimental Group A) Pre-Intervention and B) Post-Intervention (N = 7)

Variables	Groups		t-test Statistics	
	Experimental (Mean±SD)	Control (Mean±SD)	t-value	p-value
PRE-INTERVENTION				
Emotional eating scale	84.29±9.14	89.86±3.63	-1.499	0.16
PCS	47.00±6.32	51.00±11.03	-0.832	0.421
MCS	40.14±7.65	41.86±4.18	-0.52	0.612
GAD	14.00±3.37	15.71±2.14	-1.137	0.278
PHQ	18.57±4.12	18.71±2.14	-0.081	0.936
Encouragement	13.71±5.06	13.86±1.07	-0.073	0.943
Discouragement	17.57±2.07	16.00±0.00	2.008	0.068
B) POST-INTERVENTION				
Emotional Eating Scale	39.43±7.25	90.14±4.02	-16.181	0.000
PCS	67.00±1.53	47.57±15.12	3.382	0.005
MCS	59.14±1.21	38.29±7.54	7.222	0.000
GAD	0.57±0.79	16.00±1.73	-21.457	0.000
PHQ	0.71±0.76	19.00±3.70	-12.821	0.000
Encouragement	19.71±2.06	12.57±2.30	6.124	0.000
Discouragement	9.00±1.15	20.57±1.40	-16.99	0.000

Study revealed (Table 4) no significant difference for scores of EES, PCS, MCS, GAD, PHQ, encouragement and discouragement between Experimental and Control Group before intervention  $P > 0.05$ . While post-intervention results of Control versus Experimental Group as assessed using an independent sample *t*-test revealed significantly ( $p = 0.000$ ) lower EES for experimental group compared to control ( $39.43 \pm 7.25$  vs.  $90.14 \pm 4.02$ ); PCS revealed significant ( $p = .0005$ ) higher scores for Experimental Group vs Control group ( $67.00 \pm 1.53$  vs.  $47.57 \pm 15.12$ ); MCS revealed significantly ( $p = 0.000$ ) higher scores for Experimental Group vs Control ( $59.14 \pm 1.21$  vs.  $38.29 \pm 7.54$ ); GAD revealed significantly ( $p = 0.000$ ) lower scores for Experimental group vs Control ( $0.57 \pm 0.79$  vs.  $16.00 \pm 1.73$ ); PHQ revealed significantly ( $p = 0.000$ ) lower scores for experimental group compared to control ( $0.71 \pm 0.76$  vs.  $19.00 \pm 3.70$ ); Encouragement subscale revealed significantly ( $p = 0.000$ ) higher scores for Experimental group compared to control ( $19.71 \pm 2.06$  vs.  $12.57 \pm 2.30$ ) while

Discouragement subscale revealed significantly ( $p = 0.000$ ) lower scores for Experimental compared to control group ( $9.00 \pm 1.15$  vs  $20.57 \pm 1.40$ ).

These scores indicate that intervention resulted in less psychological distress as compared to control group, with higher level of encouragement and lower discouragement scores helping to control emotional eating. These findings support all four hypotheses, showing that the intervention significantly reduce emotional eating and psychological distress, increased perceived social support, and enhanced overall quality of life among overweight/obese young adults.

## Discussion

Emotional eating as a cause of weight gain cannot be ignored (Frayn & Knaüper, 2017), since literature suggests that there is association of emotional eating with weight gain (Dakanalis et al., 2023). In current study results showed higher baseline assessment score EES, GAD and discouragement. They had low social support in terms of encouragement, indicating low mental and physical health. Furthermore, the effectiveness of Intervention based on a Psycho-educational program for young adults with obesity and emotional eaters was also explored. Well reliable and valid tools were used to assess the predictors of emotional eating. Psycho-education-based intervention was based on 12 weeks, aiming to reduce BMI and emotional eating.

Literature reveals that psychological factors are involved in association of emotional dysregulation and emotional eating (De Oliveira Da Fonseca et al., 2023). In current study psycho-educational management strategy attempted to educate the university students about the results of emotional over eating, high BMI, and lower the impact of psychological issues was instituted for the interventional groups. The psycho-educational management strategy was successful in reducing emotional eating and psychological distress in terms of GAD, PHQ, and discouragement and improve the physical and mental components of QoL and encouragement among the interventional group individuals. This is in compliance with literature which suggests that management of emotional eating should target emotional regulation instead of dieting to control caloric intake (Van Strien, 2018). Association of anxiety with emotional eating was also reported in another study by Carlos et al. (2020) which recommended educational intervention. While, Tuncer and Duman (2022) reported encouraging results of using self-help program which guided the patients to regulate emotional eating. In another study Cognitive Behavioral Therapy has shown effectiveness in controlling emotional eating and overweight (Smith et

al., 2023), and a study by Eik-Nes et al. (2021), reported improvement using psychoeducational intervention for binge eating disorder with low quality of life pre intervention. Similarly, Rohrbach et al. (2022) reported internet based self-help as well as support provided to patients by clinicians are helpful in improving eating issues. According to literature based on randomized control trial study designs, psycho-educational programs have also been implemented at the junior school level, to help regulate emotions and improve problem solving skills. Significant differences were observed between baseline and post-intervention assessment (Ebrahim & Shattla, 2019). Hence, these interventions have a role to play in controlling emotional eating and weight (Treasure et al., 2022).

### **Limitations and Suggestions**

The study was conducted after the pandemic, and it is important to acknowledge that the high scores on emotional eating may have been influenced by the pandemic. Future studies should take this into account when interpreting and generalizing study findings. The current study was purely quantitative, and there may have been an impact of social desirability as participants may have felt pressure to respond in a socially desirable manner. Future studies should consider using a mixed methodology approach to gain deep knowledge from participants and gain understanding regarding major etiologies resulting in emotional eating and high BMI.

To generalize the study findings, future studies should include individuals from all age groups and population strata.

### **Conclusions**

The study found that the psycho-educational self-help weight reduction program was effective in was helpful in reducing emotional eating, psychological distress, improving social support and quality of life.

### **References**

- Carlos, M., Bernabéu, E., & Teresa, I. M. (2020). Are adherence to the Mediterranean diet, emotional eating, alcohol intake, and anxiety related in university students in Spain? *Nutrients*, *12*(8), 2224. <https://doi.org/10.3390/nu12082224>

- Coronavirus disease 2019 (COVID-19): Situation Report - 64 (24 March 2020) - World. (2020, March 24). ReliefWeb. [https://reliefweb.int/report/world/coronavirus-disease-2019-covid-19-situation-report-64-24-march-2020?gad\\_source=1&gclid=Cj0KCQiAtaOtBhCwARIsAN\\_x-3LMsqqanAAhEgQ5eekBpzfdG\\_vK5L5b7L4egu-EvnKsbyemHltfX7UaAh8GALw\\_wcB](https://reliefweb.int/report/world/coronavirus-disease-2019-covid-19-situation-report-64-24-march-2020?gad_source=1&gclid=Cj0KCQiAtaOtBhCwARIsAN_x-3LMsqqanAAhEgQ5eekBpzfdG_vK5L5b7L4egu-EvnKsbyemHltfX7UaAh8GALw_wcB).
- Dakanalis, A., Mentzelou, M., Papadopoulou, S. K., Papandreou, D., Spanoudaki, M., Vasios, G. K., Pavlidou, E., Mantzourou, M., & Giaginis, C. (2023). The Association of Emotional Eating with Overweight/Obesity, Depression, Anxiety/Stress, and Dietary Patterns: A Review of the Current Clinical Evidence. *Nutrients*, *15*(5), 1173. <https://doi.org/10.3390/nu15051173>
- Ebrahim, S. M., & Shattla, S. I. (2019). The Effectiveness of Psycho-educational Program on Emotional Regulation and Problem-Solving among Preparatory School Students. *American Journal of Nursing*, *7*(6), 922-931. <https://doi: 10.12691/ajnr-7-6-3>
- Eik-Nes, T. T., Vrabel, K., Raman, J., Clark, M. R., & Berg, K. H. (2021). A group intervention for individuals with obesity and comorbid binge eating disorder: Results from a feasibility study. *Frontiers in endocrinology*, *12*, 738856. <https://doi.org/10.3389/fendo.2021.738856>
- Emotional eating on the rise during the pandemic. (2020, November 2). MI Blues Perspectives. <https://www.mibluesperspectives.com/2020/11/02/emotional-eating-on-the-rise-during-the-pandemic/>
- Frayn, M., & Knaüper, B. (2017). Emotional Eating and Weight in Adults: a Review. *Current Psychology*, *37*(4), 924–933. <https://doi.org/10.1007/s12144-017-9577-9>
- Fonseca, N. K. D. O. D., Costa, M. D. A., Gosmann, N. P., Dalle Molle, R., Gonçalves, F. G., Silva, A. C., ... & Manfro, G. G. (2023). Emotional eating in women with generalized anxiety disorder. *Trends in Psychiatry and Psychotherapy*, *45*, e20210399. <https://doi.org/10.47626/2237-6089-2021-0399>
- Galmiche, M., Déchelotte, P., Lambert, G., & Tavolacci, M. (2019). Prevalence of eating disorders over the 2000–2018 period: a systematic literature review. *The American Journal of Clinical Nutrition*, *109*(5), 1402–1413. <https://doi.org/10.1093/ajcn/nqy342>
- Ghebreyesus, T. A. (2020). Addressing mental health needs: an integral part of COVID-19 response. *World Psychiatry*, *19*(2), 12-130. <https://doi.org/10.1002/wps.20768>
- Güneşer, R., & Him, A. (2021). Evaluation of perceived stress level and emotional eating behaviors of sports college students during COVID-19 pandemic. *Sakarya Tıp Dergisi*, *11*(2), 434-441. <https://doi.org/10.31832/smj.871642>
- Holmes, E. A., O'Connor, R. C., Perry, V. H., Tracey, I., Wessely, S., Arseneault, L., Ballard, C., Christensen, H., Silver, R. C., Everall, I., Ford, T., John, A., Kabir, T., King, K., Madan, I., Michie, S., Przybylski, A. K.,

- Shafran, R., Sweeney, A., . . . Bullmore, E. T. (2020). Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *The Lancet Psychiatry*, 7(6), 547–560. [https://doi.org/10.1016/S2215-0366\(20\)30168-1](https://doi.org/10.1016/S2215-0366(20)30168-1)
- Kroenke, K., Spitzer, R. L., & Williams, J. B. W. (2001). The PHQ-9. *Journal of General Internal Medicine*, 16(9), 606–613. <https://doi.org/10.1046/j.1525-1497.2001.016009606.x>
- Kontinen, H. (2020). Emotional eating and obesity in adults: the role of depression, sleep and genes. *Proceedings of the Nutrition Society*, 79(3), 283–289. <https://doi.org/10.1017/s0029665120000166>
- Lattimore, P. (2019). Mindfulness-based emotional eating awareness training: taking the emotional out of eating. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*, 25(3), 649-657. <https://doi.org/10.1007/s40519-019-00667-y>
- Levin, M. E., Potts, S., Haeger, J., & Lillis, J. (2018). Delivering Acceptance and Commitment Therapy for Weight Self-Stigma through Guided Self-Help: Results from an open pilot trial. *Cognitive and Behavioral Practice*, 25(1), 87–104. <https://doi.org/10.1016/j.cbpra.2017.02.002>
- McGowan, B. M. (2016). A practical guide to engaging individuals with obesity. *Obesity facts*, 9(3), 182-192. <https://doi.org/10.1159/000445193>
- Montesi, L., El Ghoch, M., Brodosi, L., Calugi, S., Marchesini, G., & Dalle Grave, R. (2016). Long-term weight loss maintenance for obesity: a multidisciplinary approach. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, 9, 37–46. <https://doi.org/10.2147/DMSO.S89836>
- Niedzwiedz, C. L., Green, M. J., Benzeval, M., Campbell, D., Craig, P., Demou, E., ... & Katikireddi, S. V. (2021). Mental health and health behaviours before and during the initial phase of the COVID-19 lockdown: longitudinal analyses of the UK Household Longitudinal Study. *J Epidemiol Community Health*, 75(3), 224-231. <https://doi.org/10.1136/jech-2020-215060>
- Rohrbach, P. J., Dingemans, A. E., Spinhoven, P., Van Ginkel, J. R., Fokkema, M., Wilderjans, T. F., Bauer, S., & Van Furth, E. F. (2022). Effectiveness of an online self-help program, expert-patient support, and their combination for eating disorders: Results from a randomized controlled trial. *International Journal of Eating Disorders*, 55(10), 1361–1373. <https://doi.org/10.1002/eat.23785>
- Sallis, J. F., Grossman, R. M., Pinski, R. B., Patterson, T. L., & Nader, P. R. (1987). The development of scales to measure social support for diet and exercise behaviors. *Preventive Medicine*, 16(6), 825–836. [https://doi.org/10.1016/0091-7435\(87\)90022-3](https://doi.org/10.1016/0091-7435(87)90022-3)
- Smith, J., Ang, X. Q., Giles, E., & Traviss-Turner, G. (2023). Emotional Eating Interventions for Adults Living with Overweight or Obesity: A Systematic Review and Meta-Analysis. *International Journal of Environmental Research and Public Health*, 20(3), 2722. <https://doi.org/10.3390/ijerph20032722>

- Spitzer, R. L., Kroenke, K., Williams, J. B. W., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder. *Archives of Internal Medicine*, *166*(10), 1092-1097. <https://doi.org/10.1001/archinte.166.10.1092>
- Tanofsky-Kraff, M., Theim, K. R., Yanovski, S. Z., Bassett, A. M., Burns, N. P., Ranzenhofer, L. M., Glasofer, D. R., & Yanovski, J. A. (2007). Validation of the emotional eating scale adapted for use in children and adolescents (EES-C). *International Journal of Eating Disorders*, *40*(3), 232–240. <https://doi.org/10.1002/eat.20362>
- Teixeira, P. J., Carraça, E. V., Marques, M. M., Rutter, H., Oppert, J. M., de Bourdeaudhuij, I., ... & Brug, J. (2015). Successful behavior change in obesity interventions in adults: a systematic review of self-regulation mediators. *BMC Medicine*, *13*(1), 84. <https://doi.org/10.1186/s12916-015-0323-6>
- Treasure, J., Bektas, S., Mutwalli, H., Dhopatkar, N., & Himmerich, H. (2022). Novel approaches to tackling emotional loss of control of eating across the weight spectrum. *Proceedings of the Nutrition Society*, *81*(3), 255–263. <https://doi.org/10.1017/s0029665122000994>
- Tuncer, G. Z., & Duman, Z. Ç. (2022). Guided self-help to manage emotional eating in a non-obese woman: A case study. *Journal of Psychiatric Nursing*, *13*(2), 184. <https://doi.org/10.14744/phd.2022.65037>
- Van Strien, T. (2018). Causes of emotional eating and matched treatment of obesity. *Current Diabetes Reports*, *18*(6), 35. <https://doi.org/10.1007/s11892-018-1000-x>
- Ware, J. E., & Sherbourne, C. D. (1992). The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *PubMed*, *30*(6), 473–483. <https://pubmed.ncbi.nlm.nih.gov/1593914>
- World Health Organization. *Obesity and Overweight*. World Health Organization: Geneva, Switzerland, 2015. URL <http://www.who.int/me-diacentre/factsheets/fs311/en/> (Accessed February 23, 2015).

Received 09 March 2024

Revision received 25 August 2025